

Selectin Biosciences Inc.

α (1-2) Fucosidase

α (1-2) Fucosidase (α -L-fucoside fucosylhydrolase, EC 3.2.1.51) cleaves non-reducing terminal fucose, linked α (1-2) to the galactose of terminal Gal-GlcNAc disaccharide structures. The presence of fucose linked to the N-acetylglucosamine will block cleavage. Substrates for α (1-2) Fucosidase are shown in Figure 1.

α (1-2) Fucosidase is isolated from *Xanthomonas*

α (1-2) Fucosidase is useful for:

- Fucose linkage determination
- Deglycosylating glycoproteins with Lewis structures

Product Code: GE 73

Specifications

Activity: ≥ 400 mU/mg, ≥ 100 mU/mL

Storage: Store at 4°C. Do not freeze.

Formulation: The enzyme is provided as a sterile-filtered solution in 20 mM Tris HCl, 25 mM NaCl pH 7.5.

Stability: Stable at least 12 months when stored properly. Several days exposure to ambient temperatures will not reduce activity.

Product Description

Molecular Weight: ~85,000 Daltons

Purity: Each lot of α (1-2) Fucosidase is tested for contaminating activities by incubating the enzyme for 24 hours at 37°C with the

appropriate substrates; the detection limit of this assay is 5 μ U/mL (IUB). A passing lot will have no detectable activity.

For the protease assay, 10 μ g of denatured BSA is incubated for 24 hours with 2 μ L of enzyme. Analysis of the BSA band after SDS-PAGE should show no evidence of degradation.

Contaminant	Substrate
β -N-acetylglucosaminidase	p-nitrophenyl- β -D-N-acetylglucosaminide
α -Galactosidase	p-nitrophenyl- α -D-galactopyranoside
β -Galactosidase	p-nitrophenyl- β -D-galactopyranoside
Neuraminidase	methylumbelliferyl- α -D-mannopyranoside
α (1-3, 4) Fucosidase	methylumbelliferyl Lewis X trisaccharide*
α -Mannosidase	p-nitrophenyl- α -D-mannopyranoside
β -Mannosidase	p-nitrophenyl- β -D-mannopyranoside
β -Xylosidase	4-methylumbelliferyl-7- α -D-xylopyranosidase

* Lewis X trisaccharide is Gal β (1-4)[Fuc α (1-3)]GlcNAc

Specificity: Non-reducing terminal fucose when linked α (1-2) to Galactose of a GalGlcNAc disaccharide structure. The presence of fucose linked to the N-acetylglucosamine will block cleavage.

Selectin Biosciences Inc.

Assay

One unit of $\alpha(1-2)$ Fucosidase is defined as the amount of enzyme required to cleave 1 μ mole of fucose from from p-nitrophenyl- α -L-fucopyranosyl- β -D-galactopyranoside in 1 minute at 37°C, pH 5.0.

Reagents

5X Reaction buffer 5.0 – 250 mM sodium phosphate, pH 5.0.

Suggestions for Use

Procedure for De-fucosylation

1. Add up to 1 nmole of oligosaccharide to tube.
2. Add deionized water to a total of 15 μ L.
3. Add 4 μ L 5X Reaction Buffer 5.0.
4. Add 1 μ L $\alpha(1-2)$ Fucosidase.
5. Incubate for 1 hour at 37°C.

Selectin Biosciences Inc warrants that the above product conforms to the specifications described herein. Should the product fail for reasons other than through misuse Selectin Biosciences Inc. will, at its option, replace free of charge or refund the purchase price. This warranty is exclusive and Selectin Biosciences Inc. makes no other warrants, expressed or implied, including any implied conditions or warranties of merchantability or fitness for any particular purpose. Selectin Biosciences Inc. shall not be liable for any incidental, consequential or contingent damages.

This product is intended for in vitro research only.

Revision 10/29/13

Selectin Biosciences Inc.

Figure 1 - Fucose residues cleaved by $\alpha(1-2)$ Fucosidase (shown in bold)

Man - Mannose; Gal - Galactose; Fuc - Fucose; GlcNAc - N-acetylglucosamine;
NeuAc - N-acetylneuraminic Acid (Sialic Acid); R - Residue

